

ASYMPTOTICS FOR FIRST-PASSAGE TIMES ON DELAUNAY TRIANGULATIONS

LEANDRO P. R. PIMENTEL

ABSTRACT. In this paper we study planar *first-passage percolation* (FPP) models, originally defined in the context of $\mathbb{Z}^d$ lattice by Hammersley and Welsh [6], on random Delaunay triangulations. The setup is as follows: to each edge $\mathbf{e}$ attach a positive random variable $\tau_{\mathbf{e}}$; the first-passage time $T(\mathbf{v}, \bar{\mathbf{v}})$ between two vertexes $\mathbf{v}$ and $\bar{\mathbf{v}}$ is defined as the infimum of $\sum_{\mathbf{e} \in \gamma} \tau_{\mathbf{e}}$ over all paths γ connecting $\mathbf{v}$ to $\bar{\mathbf{v}}$. By using subadditivity, Vahidi-Asl and Wierman [19] showed that the rescaled first-passage time converges to a constant, called the *time constant*. We show a sufficient condition to ensure that the time constant is strictly positive and derive some upper bounds for fluctuations. Our proofs are based on renormalization ideas and on the method of bounded differences.

1. INTRODUCTION

Let $\mathcal{P} \subseteq \mathbb{R}^2$ denote the set of points realized in a two-dimensional homogeneous Poisson point process with intensity 1. To each $\mathbf{v} \in \mathcal{P}$ corresponds a polygonal region $\mathbf{C}_{\mathbf{v}}$, named the *Voronoi tile* at $\mathbf{v}$, consisting of points $\mathbf{x} \in \mathbb{R}^2$ such that $|\mathbf{x} - \mathbf{v}| \leq |\mathbf{x} - \bar{\mathbf{v}}|$ for all $\bar{\mathbf{v}} \in \mathcal{P}$. The family composed by Voronoi tiles is called the *Voronoi tiling* of the plane based on $\mathcal{P}$. For a concise introduction in the subject we refer to Moller [16].

The *Delaunay Triangulation* $\mathcal{D} = (\mathcal{D}_v, \mathcal{D}_e)$ is the graph where the vertex set $\mathcal{D}_v := \mathcal{P}$ and the edge set $\mathcal{D}_e$ consists of non-oriented pairs $(\mathbf{v}, \bar{\mathbf{v}})$ such that $\mathbf{C}_{\mathbf{v}}$ and $\mathbf{C}_{\bar{\mathbf{v}}}$ share a one-dimensional boundary (Figure 1). One can see that (with probability one) each Voronoi tile is a convex and bounded polygon, and the graph $\mathcal{D}$ is a triangulation of the plane.




FIGURE 1. The Voronoi Tiling and Delaunay Triangulation.

The *Voronoi Tessellation* $\mathcal{V} = (\mathcal{V}_v, \mathcal{V}_e)$ is the graph where the vertex set $\mathcal{V}_v$ consists of vertexes of the Voronoi tiles and the edge set $\mathcal{V}_e$ is the set of edges of the Voronoi tiles. The edges $\mathbf{e}^*$ of $\mathcal{V}$ are segments of the perpendicular bisectors of the edges $\mathbf{e}$ of $\mathcal{D}$. This establishes duality of $\mathcal{D}$ and $\mathcal{V}$ as planar graphs.

To each edge $\mathbf{e} \in \mathcal{D}_e$ is independently assigned a nonnegative random variable $\tau_{\mathbf{e}}$ from a common distribution $\mathbb{F}$, which is also independent of the Poisson point process that generates $\mathcal{P}$. We assume that both $\mathcal{P}$ and $\{\tau_{\mathbf{e}} : \mathbf{e} \in \mathcal{D}_e\}$ are functions of a configuration $\omega \in \Omega$ and denote by $\mathbb{P}$ its joint law. The expectation and the variance are denoted by $\mathbb{E}$ and by $\mathbb{V}$, respectively.

Let

$$t(\gamma) := \sum_{\mathbf{e} \in \gamma} \tau_{\mathbf{e}},$$

be the *passage time* along the path γ . The *first-passage time* between two vertexes $\mathbf{v}$ and $\mathbf{v}'$ is defined by

$$T(\mathbf{v}, \mathbf{v}') := \inf\{t(\gamma); \gamma \in \mathcal{C}(\mathbf{v}, \mathbf{v}')\}$$

where $\mathcal{C}(\mathbf{v}, \mathbf{v}')$ denotes the set of all self-avoiding paths connecting $\mathbf{v}$ to $\mathbf{v}'$. A *geodesic* connecting $\mathbf{v}$ to $\mathbf{v}'$ is a path $\rho(\mathbf{v}, \mathbf{v}') \in \mathcal{C}(\mathbf{v}, \mathbf{v}')$ which attains the minimum:

$$t(\rho(\mathbf{v}, \mathbf{v}')) = T(\mathbf{v}, \mathbf{v}')$$

(a sufficient condition on $\mathbb{F}$ to ensure the existence of geodesics is given by Corollary 3.2).

For each $\mathbf{x} \in \mathbb{R}^2$ we denote $\mathbf{v}(\mathbf{x})$ the almost-surely unique point $\mathbf{v} \in \mathcal{P}$ such that $\mathbf{x} \in \mathbf{C}_{\mathbf{v}}$. For $\mathbf{x}, \mathbf{y} \in \mathbb{R}^2$ let

$$T(\mathbf{x}, \mathbf{y}) := T(\mathbf{v}(\mathbf{x}), \mathbf{v}(\mathbf{y}))$$

and

$$\rho(\mathbf{x}, \mathbf{y}) := \rho(\mathbf{v}(\mathbf{x}), \mathbf{v}(\mathbf{y})).$$

The set of points reached from $\mathbf{x}$ by time t is defined by

$$\mathbf{B}_{\mathbf{x}}(t) := \{\mathbf{y} \in \mathbb{R}^2 : \mathbf{y} \in \mathbf{C}_{\mathbf{v}} \text{ where } \mathbf{v} \in \mathcal{D}_v \text{ and } T(\mathbf{v}(\mathbf{x}), \mathbf{v}) \leq t\},$$

We remark that others FPP models (euclidean FPP) were introduced by Howard and Newman [7], where the underline graph is the complete graph with vertex set $\mathcal{P}$ and to each edge $\mathbf{e} = (\mathbf{v}, \bar{\mathbf{v}})$ is attached the passage time $\tau_{(\mathbf{v}, \bar{\mathbf{v}})} := |\mathbf{v} - \bar{\mathbf{v}}|^\alpha$ ($\alpha > 0$ is a fixed parameter).

We recall fundamental results in the subjects, proved by Vahidi-Asl and Wierman [19, 20]. Let

$$\mu(\mathbb{F}) := \inf_{n>0} \frac{\mathbb{E}T(\mathbf{0}, \mathbf{n})}{n} \in [0, \infty]$$

where $\mathbf{0} := (0, 0)$ and $\mathbf{n} := (n, 0)$. $\mu(\mathbb{F})$ is usually refereed the *time constant*. Assume that τ_1, τ_2, τ_3 are independent random variables with common distribution $\mathbb{F}$: if

$$\mathbb{E}(\min_{j=1,2,3} \{\tau_j\}) < \infty \tag{1.1}$$

then $\mu(\mathbb{F}) < \infty$ and for all unit vectors $\vec{\mathbf{x}} \in S^1$ ($|\vec{\mathbf{x}}| = 1$) $\mathbb{P}$ -a.s.

$$\lim_{n \rightarrow \infty} \frac{T(\mathbf{0}, n\vec{\mathbf{x}})}{n} = \lim_{n \rightarrow \infty} \frac{\mathbb{E}T(\mathbf{0}, \mathbf{n})}{n} = \mu(\mathbb{F}). \quad (1.2)$$

In addition, if

$$\mathbb{E}(\min_{j=1,2,3} \{\tau_j\}^2) < \infty$$

holds and $\mu(\mathbb{F}) > 0$ then for all $\epsilon > 0$ $\mathbb{P}$ -a.s. there exists $t_0 > 0$ such that for all $t > t_0$

$$(1 - \epsilon)t\mathbf{D}(1/\mu) \subseteq \mathbf{B}_0(t) \subseteq (1 + \epsilon)t\mathbf{D}(1/\mu),$$

where $\mathbf{D}(r) := \{\mathbf{x} \in \mathbb{R}^2 : |\mathbf{x}| \leq r\}$.

Two natural questions arise from (1.2).

- When is $\mu(\mathbb{F}) > 0$?
- What is the right order of $T(\mathbf{0}, \mathbf{n}) - \mu n$?

It is expected that

$$\mathbb{F}(0) < p_c := \inf\{p > 0; \theta(p) = 1\},$$

where $\theta(p)$ is the probability that bond percolation on $\mathcal{D}$ occurs with density p , is a sufficient and necessary condition to have $\mu(\mathbb{F}) > 0$ (Kesten [12]).

Heuristics arguments indicate that FPP models belongs to the KPZ universality class, which brings the conjecture that the right order is n^χ , where $\chi = 1/3$ (Kadar, Parisi and Zhang [10], Krug and Spohn [14]). However the only models for which this has been proved are certain growth models related to random permutations (Baik, Deift and Johansson [2], Johansson [9]). For lattice FPP Kesten [13] showed that $\chi \leq 1/2$ (see also Alexander [1]), and for euclidean FPP Howard and Newman [8] showed the same upper bound.

In this work we prove some results related to the questions above. To state them we require some definitions involving a bond percolation model.

Let $\mathbf{A}$ and $\mathbf{B}$ be two subsets of $\mathbb{R}^2$. Let $[\mathbf{x}, \mathbf{y}]$ denote the line segment connecting $\mathbf{x}$ to $\mathbf{y}$. We say that a self-avoiding path $\gamma^* = (\mathbf{v}_1^*, \dots, \mathbf{v}_k^*)$ in $\mathcal{V}$ is a path connecting $\mathbf{A}$ to $\mathbf{B}$ if $[\mathbf{v}_1^*, \mathbf{v}_2^*] \cap \mathbf{A} \neq \emptyset$ and $[\mathbf{v}_{k-1}^*, \mathbf{v}_k^*] \cap \mathbf{B} \neq \emptyset$. The bond percolation model on the Voronoi Tessellation $\mathcal{V}$, with parameter p and probability law $\mathbb{P}_p^*$, is constructed by choosing each edge $\mathbf{e}^*$ in $\mathcal{V}_e$ to be open independently with probability p . An open path is a path composed by open edges. For each $R > 0$ denote by A_R the event that there exists an open path $\gamma = (\mathbf{v}_j^*)_{1 \leq j \leq h}$ in $\mathcal{V}$, connecting $\{0\} \times [0, R]$ to $\{3R\} \times [0, R]$, and with $\mathbf{v}_j^* \in [0, 3R] \times [0, R]$ for all $j = 2, \dots, h-1$. Define the function

$$\eta^*(p) := \liminf_{R \rightarrow \infty} \mathbb{P}_p^*(A_R),$$

and consider the percolation threshold,

$$p_c^* := \inf\{p > 0 : \eta^*(p) = 1\} < 1 \quad (1.3)$$

(the inequality in (1.3) will follow from Proposition 3.1).

For each $\kappa > 1/2$ we set $\nu = \nu(\kappa) := (4\kappa - 2)/7$. For the remainder of the paper we will use the symbols a_j, c_j, b_j for $j = 1, 2, \dots$ to represent strictly positive constants, whose value may change from appearance to appearance but will not depend on n, r, s or t in the notation we will follow here.

Theorem 1. *If $\mathbb{F}(0) < 1 - p_c^*$ and (1.1) holds then*

$$0 < \mu(\mathbb{F}) < \infty. \quad (1.4)$$

If (1.1) is strengthened to

$$\mathbb{E}(e^{a\tau}) = \int e^{at} d\mathbb{F}(t) < \infty \text{ for some } a > 0 \quad (1.5)$$

then:

- *For all $\epsilon > 0$ there exists $n_0 > 0$ such that for all $n > n_0$*

$$\mathbb{V}(T(\mathbf{0}, \mathbf{n})) \leq n^{1+\epsilon}; \quad (1.6)$$

- *For all $\kappa > 1/2$ there exist constants $c_j > 0$ such that for all $n \geq 1$ and $r \in [0, c_1 n^\kappa]$*

$$\mathbb{P}(|T(\mathbf{0}, \mathbf{n}) - \mathbb{E}T(\mathbf{0}, \mathbf{n})| \geq rn^\kappa) \leq e^{-c_2 r^\nu}, \quad (1.7)$$

and

$$\mu n \leq \mathbb{E}T(\mathbf{0}, \mathbf{n}) \leq \mu n + c_3 n^\kappa (\log n)^{1/\nu}. \quad (1.8)$$

We note that our condition to get $\mu(\mathbb{F}) > 0$ should be equivalent to $\mathbb{F}(0) < p_c$, since it is expected that $p_c + p_c^* = 1$ (duality) for many planar graphs¹.

From Theorem 1, the proof of which is given in Section 4, we get an upper bound for the fluctuations of $T(\mathbf{0}, \mathbf{n})$ about μn and an improved shape theorem.

Corollary 1.1. *If $\mathbb{F}(0) < 1 - p_c^*$ and (1.5) holds then for all $\kappa > 1/2$ there exist constants $c_j > 0$ such that for all $r \in [c_3(\log n)^{1/\nu}, c_1 n^\kappa]$*

$$\mathbb{P}(|T(\mathbf{0}, \mathbf{n}) - \mu n| \geq rn^\kappa) \leq e^{-\frac{c_2}{2} r^\nu}. \quad (1.9)$$

Proof of Corollary 1.1. (1.9) follows immediately from (1.7) and (1.8). □

Corollary 1.2. *If $\mathbb{F}(0) < 1 - p_c^*$ and (1.5) holds then for all $\kappa > 1/2$, $\mathbb{P}$ -a.s. there exists $t_0 > 0$ such that for all $t > t_0$*

$$(t - t^\kappa)\mathbf{D}(1/\mu) \subseteq \mathbf{B}_0(t) \subseteq (t + t^\kappa)\mathbf{D}(1/\mu).$$

¹It follows from (1.4) that $p_c + p_c^* \geq 1$ holds (Pimentel [17])

Proof of Corollary 1.2. Notice that, for large $|\mathbf{x}|$, if

$$|T(\mathbf{0}, \mathbf{x}) - \mu|\mathbf{x}|| > |\mathbf{x}|^\kappa$$

then

$$|T(\mathbf{0}, \mathbf{z}) - \mu|\mathbf{z}|| > |\mathbf{z}|^\kappa/4 \text{ or } \max_{|\mathbf{x}-\mathbf{z}|\leq 1} T(\mathbf{x}, \mathbf{z}) > |\mathbf{z}|^\kappa/4. \quad (1.10)$$

Under (1.5), one can easily see that $\mathbb{P}$ -a.s. the event in the right hand side of (1.10) occurs for finitely many $\mathbf{z} \in \mathbb{Z}^2$. From (1.9) (and the Borel-Cantelli lemma) it follows that $\mathbb{P}$ -a.s. the event in the left-hand side of (1.10) occurs for finitely many $\mathbf{z} \in \mathbb{Z}^2$.

Therefore, $\mathbb{P}$ -a.s. there exist $M > 0$ such that for all $\mathbf{x} \in \mathbb{R}^2$ with $|\mathbf{x}| > M$

$$|T(\mathbf{0}, \mathbf{x}) - \mu|\mathbf{x}|| \leq |\mathbf{x}|^\kappa,$$

which yields Corollary 1.2 (further details are left to the reader). □

Overview. In Section 2 we will study some geometrical aspects of Voronoi tilings and of self-avoiding paths on Delaunay triangulations, which will play an important role in control the asymptotic behavior of first-passage times. In Section 3 we explore the duality between bond percolation models on Delaunay triangulations and on Voronoi tessellations to relate the size of $\mathbb{F}(0)$ and the value of $\mu(\mathbb{F})$. The fluctuations of the first-passage time about its asymptotic value are considered in Section 4.

2. PRELIMINARIES

The main tool we handle to control the fluctuations of the first-passage time about its asymptotic value is the *method of bounded increments*, which was also considered by Kesten [13] in the lattice FPP context, and also by Howard and Newman [8] in the euclidean FPP context, to study the same question. This method represents $T(\mathbf{0}, \mathbf{n}) - \mathbb{E}T(\mathbf{0}, \mathbf{n})$ as a sum of martingales increments and, after estimating these increments, applies standard bounds for martingales with bounded increments.

One of the great difficulties here is that, as distinct from the lattice context, a local increment in a configuration ω is felt not only by the travel times but also by the (random) graph, which has a long range dependence. For this reason we shall define a truncation of the Poisson process.

For $\mathbf{z} \in \mathbb{Z}^2$, $r > 0$ and $s \in \{j/2 : j \in \mathbb{N}\}$ let

$$\mathbf{B}_{\mathbf{z}}^{s,r} := r\mathbf{z} + [-sr, sr]^2. \quad (2.11)$$

Order the points of $\mathbb{Z}^2$ in some arbitrary fashion, say $\mathbb{Z}^2 := \{\mathbf{u}_1, \mathbf{u}_2, \dots\}$. Let $\delta > 0$ be a fixed parameter which value will be specified later ². Let $n \geq 1$ and for each $k \geq 1$ let

$$\mathbf{B}_k^n := \mathbf{B}_{\mathbf{u}_k}^{1/2, n^\delta}.$$

²Its value will depend on ϵ and κ in Theorem 1

Let $|\mathbf{A}|$ denote the number of elements belonging to the set $\mathbf{A}$. Define the point process $\mathcal{P}_n := \mathcal{P}_n(\mathcal{P})$ (whose distribution will also depend on δ) as follows:

- If $|\mathbf{B}_k^n \cap \mathcal{P}| = 0$ then set $\mathbf{B}_k^n \cap \mathcal{P}_n := \{\mathbf{v}_k\}$ where $\mathbf{v}_k$ is a random point uniformly distributed in $\mathbf{B}_k^n$;
- If $|\mathbf{B}_k^n \cap \mathcal{P}| > 4n^{2\delta}$ then set $\mathbf{B}_k^n \cap \mathcal{P}_n := \{\mathbf{v}_k^1, \dots, \mathbf{v}_k^{4n^{2\delta}}\}$ where $(\mathbf{v}_k^1, \dots, \mathbf{v}_k^{4n^{2\delta}})$ is a sequence of $4n^{2\delta}$ points uniformly chosen in $\mathbf{B}_k^n \cap \mathcal{P}$.

We make the convention $\mathcal{P}_\infty = \mathcal{P}$ and denote by $\mathcal{D}_n$ the Delaunay Triangulation based on $\mathcal{P}_n$.

2.1. Renormalization and full boxes. The geometry of Voronoi tilings is study through renormalization ideas. In a few words, it consists in defining a larger box to be a *good box* if some property is satisfied in a neighborhood of that box. In the course of the proofs we shall utilize different definitions of good boxes but all of them will carry the notion of *full box*.

Formally, we divide a square box $\mathbf{B} = \mathbf{B}_{\mathbf{z}}^{1/2,r}$ into thirty-six sub boxes of the same length, say $\mathbf{B}_1, \dots, \mathbf{B}_{36}$. We stipulate $\mathbf{B}$ is a full box, with respect to the point configuration $\mathcal{P}_n$, if all those thirty-six sub boxes have at least one point belonging to $\mathcal{P}_n$ (Figure 2). In other words,

$$[\mathbf{B} = \cup_{j=1}^{36} \mathbf{B}_j \text{ is a full box}] := [\mathbf{B}_j \cap \mathcal{P}_n \neq \emptyset \forall j = 1, \dots, 36]. \quad (2.12)$$




FIGURE 2. Renormalization: a full box

We say that $\Lambda := (\mathbf{B}_{\mathbf{z}_1}^{1/2,r}, \dots, \mathbf{B}_{\mathbf{z}_k}^{1/2,r})$ is a circuit of boxes if $(\mathbf{z}_1, \dots, \mathbf{z}_k)$ is a circuit in $\mathbb{Z}^2$. Let λ be the closed polygonal path composed by the line segments connecting $r\mathbf{z}_j$ to $r\mathbf{z}_{j+1}$, where $j = 1, \dots, k-1$, together with $[\mathbf{z}_k, \mathbf{z}_1]$. To each circuit Λ we associate four subsets of the plane: Λ^{out} will denote the (topological) interior of the unbounded component of $\mathbb{R}^2 \setminus \cup_{j=1}^k \mathbf{B}_{\mathbf{z}_j}^{1/2,r}$, while Λ^{in} will denote the interior of the bounded component of $\mathbb{R}^2 \setminus \cup_{j=1}^k \mathbf{B}_{\mathbf{z}_j}^{1/2,r}$; λ^{out} will denote the interior of the unbounded component of $\mathbb{R}^2 \setminus \lambda$, while λ^{in} will denote the interior of the bounded component of $\mathbb{R}^2 \setminus \lambda$.

Lemma 2.1. *Assume that $\Lambda := (\mathbf{B}_{\mathbf{z}_j}^{1/2,r})_{j=1}^k$ is a circuit composed by full boxes with respect to $\mathcal{P}_n$.*

$$\text{If } \mathbf{C}_{\mathbf{v}} \cap \Lambda^{in} \neq \emptyset \Rightarrow \mathbf{C}_{\mathbf{v}} \subseteq \lambda^{in}. \quad (2.13)$$

$$\text{If } \mathbf{C}_{\mathbf{v}} \cap \Lambda^{out} \neq \emptyset \Rightarrow \mathbf{C}_{\mathbf{v}} \subseteq \lambda^{out}. \quad (2.14)$$

Proof of Lemma 2.1. By convexity of Voronoi tiles, if (2.13) does not hold then there exist $\mathbf{x}_1 \in \partial\Lambda^{in} \cap \mathbf{C}_{\mathbf{v}}$ and $\mathbf{x}_2 \in \partial\Lambda^{in} \cap \mathbf{C}_{\mathbf{v}}$ ($\partial\mathbf{A}$ denotes the boundary of the set $\mathbf{A}$) and thus

$$r/2 \leq |\mathbf{x}_1 - \mathbf{x}_2|.$$

Since every box in $(\mathbf{B}_{\mathbf{z}_j}^{1/2,r})_{j=1}^k$ is a full box, there exist $\mathbf{v}_1, \mathbf{v}_2 \in \mathcal{P}_n$ so that

$$|\mathbf{v}_1 - \mathbf{x}_1| \leq \sqrt{2}r/6 \text{ and } |\mathbf{v}_2 - \mathbf{x}_2| \leq \sqrt{2}r/6.$$

Although, $\mathbf{x}_1$ and $\mathbf{x}_2$ belong to $\mathbf{C}_{\mathbf{v}}$ and so

$$|\mathbf{v} - \mathbf{x}_1| \leq |\mathbf{v}_1 - \mathbf{x}_1| \text{ and } |\mathbf{v} - \mathbf{x}_2| \leq |\mathbf{v}_2 - \mathbf{x}_2|.$$

Thus,

$$r/2 \leq |\mathbf{x}_1 - \mathbf{x}_2| \leq |\mathbf{x}_1 - \mathbf{v}| + |\mathbf{x}_2 - \mathbf{v}| \leq \sqrt{2}r/3,$$

which leads to a contradiction since $\sqrt{2}/3 < 1/2$. By an analogous argument, one can prove (2.14). □

For each $\mathbf{A} \subseteq \mathbb{R}^2$ let $\mathcal{G}_{\mathcal{P}_n}(\mathbf{A})$ be the sub-graph of $\mathcal{D}_n$ composed by edges $\mathbf{e} = (\mathbf{v}_1, \mathbf{v}_2)$ so that $\mathbf{C}_{\mathbf{v}_i} \cap \mathbf{A} \neq \emptyset$ for $i = 1, 2$. We denote by $\tau(\mathbf{A})$ the sequence of passage times associated to edges in $\mathcal{G}_{\mathcal{P}_n}(\mathbf{A})$.

Given a random variable X and a measurable event F we denote by $X \mid F$ the random variable X condition on the event F .

Lemma 2.2. *Let F_Λ be the event that $\Lambda := (\mathbf{B}_{\mathbf{z}_j}^{1/2,r})_{j=1}^k$ is a circuit composed by full boxes with respect to $\mathcal{P}_n$. Assume that X is a random variable which only depend on $(\mathcal{G}_{\mathcal{P}_n}(\Lambda^{in}), \tau(\Lambda^{in}))$ and that Y is a random variable which only depend on $(\mathcal{G}_{\mathcal{P}_n}(\Lambda^{out}), \tau(\Lambda^{out}))$. Then*

$$X \mid F_\Lambda \text{ is independent of } Y \mid F_\Lambda.$$

Proof of Lemma 2.2. It follows from Lemma 2.1. □

2.2. Greedy lattice animals and site percolation. In this section we introduce two models which will also play an important role in the study of Voronoi tilings. The first one, named *Greedy lattice animals* and introduced by Cox, Gandolfi, Griffin and Kesten [3]), consists of the following: a lattice animal is a connected subset of $\mathbb{Z}^2$ containing the origin. Let Φ^r and Φ_r denote the set of animals with at most r sites and with at least r sites, respectively.

Let $X := \{X_{\mathbf{z}} : \mathbf{z} \in \mathbb{Z}^2\}$ be a collection of i.i.d. non-negative random variables and define

$$M_s(X) := \max\left\{\sum_{\mathbf{z} \in \mathbf{A}} X_{\mathbf{z}} : \mathbf{A} \in \Phi^s\right\}. \quad (2.15)$$

A *greedy lattice animal* is a lattice animal which attains the maximum in the above definition.

Lemma 2.3. *If*

$$\mathbb{E}(e^{cX_0}) < \infty \text{ for some } c > 0 \quad (2.16)$$

holds then there exist $b_1, b_2 > 0$ such that

$$\mathbb{P}(M_s > b_1 s) \leq e^{-b_2 s}.$$

Proof of Lemma 2.3. Under (2.15),

$$\mathbb{E}(M_s) \leq bs$$

for some $b > 0$ (Cox, Gandolfi, Griffin, Kesten [3]). From Corollary 8.2.4 of Talagrand [18], if v_s denotes a median of M_s then

$$\mathbb{P}(|M_s - v_s| > u) \leq 4 \exp(-k_0 \min\{s^{-1}u^2, u\}).$$

for some constant $c_0 > 0$.

By integrating in u from 0 to ∞ both sides of the last inequality we get that, for some constant $\bar{b} > 0$,

$$|\mathbb{E}M_s - v_s| \leq \mathbb{E}|M_s - v_s| \leq \bar{b}\sqrt{s}.$$

and so

$$\mathbb{P}(|M_s - \mathbb{E}M_s| > u) \leq \exp(-c_0 \min\{s^{-1}u^2, u\}),$$

if $u > 2\bar{b}\sqrt{s}$. This implies that

$$\mathbb{P}(M_s \geq 2bs) \leq \mathbb{P}(|M_s - \mathbb{E}M_s| > bs) \leq e^{-bs/2c_0}, \quad (2.17)$$

which yields Lemma 2.3. □

Now, for each $n \geq 1$ consider $Y^n := \{Y_{\mathbf{z}}^n : \mathbf{z} \in \mathbb{Z}^2\}$, a collection of Bernoulli random variables. Let

$$\phi := \inf_{n \geq 1} \inf_{\mathbf{z} \in \mathbb{Z}^2} \mathbb{P}(Y_{\mathbf{z}}^n = 1).$$

Assume that this collection is l -dependent: $\{Y_{\mathbf{z}}^n : \mathbf{z} \in \mathbf{A}\}$ and $\{Y_{\mathbf{z}}^n : \mathbf{z} \in \mathbf{B}\}$ are independent whenever

$$l \leq d_\infty(\mathbf{A}, \mathbf{B}) := \min\{|\mathbf{z} - \mathbf{z}'|_\infty : \mathbf{z} \in \mathbf{A} \text{ and } \mathbf{z}' \in \mathbf{B}\}.$$

This defines a family of l -dependent site percolation models index by $n \in [1, \infty]$ (a site $\mathbf{z}$ is called *open* if $Y_{\mathbf{z}}^n = 1$). Here we are interested on the density of open sites in the set of lattice animals with at least s sites. For more details in the subject see the reference book of Grimmet [4].

For each animal $\mathbf{A}$ we denote by $\Xi_{\mathbf{A}}^k$ the set of all $\mathbf{B} \subseteq \mathbf{A}$ such that if $\mathbf{z}, \bar{\mathbf{z}} \in \mathbf{B}$ then $d_\infty(\mathbf{z}, \bar{\mathbf{z}}) \geq k$. Let

$$m^k(\mathbf{A}, Y^n) := \max\left\{\sum_{\mathbf{z} \in \mathbf{B}} Y_{\mathbf{z}}^n : \mathbf{B} \in \Xi_{\mathbf{A}}^k\right\}$$

and

$$m_s^k(Y^n) := \min\{m^k(\mathbf{A}, Y^n) : \mathbf{A} \in \Phi_s\}. \quad (2.18)$$

Lemma 2.4. *For all $l \geq 1$ there exists $p_0 \in (0, 1)$ such that if $\phi > p_0$ then for all $k \geq 1$ there exist $b_3, b_4 > 0$ such that for all $n \in [1, \infty]$ and $s \geq 1$,*

$$\mathbb{P}(m_s^k(Y^n) < b_3 s) \leq e^{-b_4 s}. \quad (2.19)$$

Proof of Lemma 2.4. By Theorem 0.0 of Liggett, Schomman, Stacey [15], the family

$$\{Y^n : n \in [1, \infty]\}$$

is stochastically dominated from below by a collection of i.i.d Bernoulli random variables with parameter $\rho(\phi) \rightarrow 1$ when $\phi \rightarrow 1$.

Therefore, Lemma 2.4 will follow if we prove that (2.19) holds for $m_s^k(Y)$, where $Y := \{Y_{\mathbf{z}} : \mathbf{z} \in \mathbb{Z}^2\}$ is a collection of i.i.d Bernoulli random variables with parameter ρ sufficiently close to 1. Since $m_s^1(Y) \leq c(k)m_s^k(Y)$, for some constant $c(k)$ (only depending on k), we may restrict our attention to $k = 1$.

For each animal $\mathbf{A}$, $\sum_{\mathbf{z} \in \mathbf{A}} Y_{\mathbf{z}}$ is a binomial random variable with parameter $|\mathbf{A}|$ and ρ . Since (the number of animals $\mathbf{A}$ with $|\mathbf{A}| = j$ is at most C^j , for some constant $C > 0$) we have at most c^j animals with $|\mathbf{A}| = j$, we get that

$$\begin{aligned} \mathbb{P}(m_s(Y)^1 < bs) &= \mathbb{P}(\exists \mathbf{A} : \sum_{\mathbf{z} \in \mathbf{A}} Y_{\mathbf{z}} < bs \text{ and } |\mathbf{A}| \geq s) \\ &\leq \sum_{j \geq s} c^j \sum_{l=0}^{bs} C_l^j \rho^l (1-\rho)^{j-l} \leq (1-\rho)^{-bs} \sum_{j \geq s} [2c(1-\rho)]^j \\ &= \frac{(1-\rho)^{-bs} [2c(1-\rho)]^s}{1 - 2c(1-\rho)} = \frac{\exp\{s \log 2c + s(1-b) \log(1-\rho)\}}{1 - 2c(1-\rho)} \end{aligned}$$

if $b < 1$ and $\rho > 1 - 1/2c$. Thus, if we fix $\rho > 1 - 1/2c$ and

$$b_3 < 1 + \log 2c [\log(1-\rho)]^{-1}$$

then we can find $b_4 = b_4(\rho) > 0$ so that (2.19) holds for $m_s^1(Y)$.

□

2.3. Self-avoiding paths on Delaunay triangulations. Let $\mathcal{C}_r$ and $\mathcal{C}^r$ denote the set of self-avoiding and finite paths γ in $\mathcal{D}_n$, starting from $\mathbf{v}(\mathbf{0})$, with $|\gamma| \geq r$ and with $|\gamma| \leq r$, respectively.

Fix $L > 0$ and for each path γ in $\mathcal{D}_n$ let $\mathbf{A}(\gamma) \subseteq \mathbb{Z}^2$ be the set of points $\mathbf{z} \in \mathbb{Z}^2$ so that $\mathbf{B}_{\mathbf{z}}^{1/2, L} \cap [\mathbf{v}, \mathbf{v}'] \neq \emptyset$ for some $\mathbf{v}, \mathbf{v}' \in \gamma$ (Figure 3). Define

$$g_r^L(\mathcal{P}_n) := \min\{|\mathbf{A}(\gamma)| : \gamma \in \mathcal{C}_r(\mathcal{P}_n)\}$$

and

$$G_r^L(\mathcal{P}_n) := \max\{|\mathbf{A}(\gamma)| : \gamma \in \mathcal{C}^r(\mathcal{P}_n)\}.$$




FIGURE 3. γ and $\mathbf{A}(\gamma)$

Let $\mathbf{x}, \mathbf{y} \in \mathbb{R}^2$ and define the path $\gamma(\mathbf{x}, \mathbf{y}) := (\mathbf{v}_1, \dots, \mathbf{v}_k)$ in $\mathcal{D}(\mathcal{P}_n)$ as follows: set $\mathbf{v}_1 := \mathbf{v}(\mathbf{x})$; if $\mathbf{v}_1 \neq \mathbf{v}(\mathbf{y})$ then let $\mathbf{v}_2$ be the (almost-surely) unique nearest neighbor of $\mathbf{v}_1$ such that the edge of $\mathbf{C}_{\mathbf{v}_1}$, that is perpendicular to $[\mathbf{v}_1, \mathbf{v}_2]$, crosses $[\mathbf{x}, \mathbf{y}]$; otherwise we set $k = 1$ and the construction is finished; given $\mathbf{v}_l$ with $l \geq 1$, if $\mathbf{v}_l \neq \mathbf{v}(\mathbf{y})$ then we set $\mathbf{v}_{l+1}$ to be the unique nearest neighbor of $\mathbf{v}_l$ different from $\mathbf{v}_{l-1}$ such that the edge of $\mathbf{C}_{\mathbf{v}_l}$, that is perpendicular to $[\mathbf{v}_l, \mathbf{v}_{l+1}]$, crosses $[\mathbf{x}, \mathbf{y}]$; otherwise we set $k := l$ and the construction is finished. We denote $\gamma_r := \gamma(\mathbf{0}, r\mathbf{1})$.

Proposition 2.1. *For each $L > 1$ there exist $c_j = c_j(L) > 0$ such that for all $n \in [1, \infty]$, for all $r > 0$,*

$$\mathbb{P}(g_r^L(\mathcal{P}_n) < c_1 r) \leq e^{-c_2 r}, \quad (2.20)$$

and

$$\mathbb{P}(G_r^L(\mathcal{P}_n) > c_3 r) \leq e^{-c_4 r}. \quad (2.21)$$

Proposition 2.2. *There exists $z_0 > 0$ such that, for all $n \in [1, \infty]$, for all $r > 0$ and $z > z_0$*

$$\mathbb{P}(|\gamma_r(\mathcal{P}_n)| > zr) \leq e^{-c_1 z r}. \quad (2.22)$$

Proof of Proposition 2.1. First notice that to prove (2.20) we can restrict the attention to $L = 1$. Let $X_{\mathbf{z}}^n := |\mathbf{B}_{\mathbf{z}}^{1/2,1} \cap \mathcal{P}_n|$ and denote $X^n := \{X_{\mathbf{z}}^n : \mathbf{z} \in \mathbb{Z}^2\}$. Consider the random variable $M_s^n := M_s(X^n)$ defined in (2.15).

By definition, if $g_r^1(\mathcal{P}_n) \leq cr$ then there exists a path $\gamma \in \mathcal{C}_r(\mathcal{P}_n)$ with $|A(\gamma)| \leq cr$. Thus

$$M_{cr}^n \geq \sum_{z \in \mathbf{A}(\gamma)} |\mathbf{B}_z^{1/2,1} \cap \mathcal{P}_n| \geq |\gamma| \geq r,$$

which yields that

$$\mathbb{P}(g_r^1(\mathcal{P}_n) \leq cr) \leq \mathbb{P}(M_{cr}^n \geq r). \quad (2.23)$$

By the definition of $\mathcal{P}_n$, for each $\mathbf{z} \in \mathbb{Z}^2$

$$|\mathbf{B}_{\mathbf{z}}^{1/2,1} \cap \mathcal{P}_n| \leq |\mathbf{B}_{\mathbf{z}}^{1/2,1} \cap \mathcal{P}_\infty| + 1,$$

and then

$$M_r^n \leq M_r^\infty + r.$$

Combining this with (2.23), one gets that

$$\mathbb{P}(g_r^1(\mathcal{P}_n) \leq cr) \leq \mathbb{P}(M_{cr}^\infty \geq (1-c)r).$$

Since (2.16) holds for X^∞ , together with Lemma 2.3, this implies (2.20).

The proof of (2.21) is founded on renormalization techniques as follows. Denote $\mathbf{C}_{\mathbf{z}}$ the circuit composed by sites $\mathbf{z}' \in \mathbb{Z}^2$ with $|\mathbf{z} - \mathbf{z}'|_\infty = 1$. We say that $\mathbf{B}_{\mathbf{z}}^{1/2,L}$ is good box, or equivalently $Y_{\mathbf{z}}^n(L) = 1$, if $\mathbf{B}_{\mathbf{z}}^{1/2,L}$ is a full box with respect to $\mathcal{P}_n$ for all $\bar{\mathbf{z}} \in \mathbf{C}_{\mathbf{z}}$ (see (2.12)). We denote $Y^n(L) := \{Y_{\mathbf{z}}^n(L) : \mathbf{z} \in \mathbb{Z}^2\}$.

Lemma 2.5. *$\{Y_{\mathbf{z}}^n(L) : \mathbf{z} \in \mathbb{Z}^2\}$ is a 3-dependent collection of Bernoulli random variables. Further, for all $\epsilon > 0$ there exists $L_0 > 0$ so that*

$$\inf_{n \geq 1} \inf_{\mathbf{z} \in \mathbb{Z}^2} \mathbb{P}(Y_{\mathbf{z}}^n(L_0)) \geq 1 - \epsilon.$$

Proof of Lemma 2.5. Since $\mathbf{C}_z$ and $\mathbf{C}_{z'}$ are disjoint if $|\mathbf{z} - \mathbf{z}'| \geq 3$, the first part of this lemma follows directly from the definition of $\mathcal{P}_n$.

To prove the second part notice that if $L/36 \geq n^\delta$ then each sub-box of $\mathbf{B}_{\mathbf{z}}^{1/2,L}$ with length $L/6$ contains at least one box $\mathbf{B}_n^k$ (used to construct $\mathcal{P}_n$). Thus

$$\mathbb{P}(Y_{\mathbf{z}}^L(\mathcal{P}_n) = 1) = 1.$$

Now, if $L/36 < n^\delta$ and $\mathbf{B}$ is a box with length $L/6$ then

$$\begin{aligned} \mathbb{P}(\mathbf{B} \cap \mathcal{P}_n \neq \emptyset) &\leq \mathbb{P}(\mathbf{B} \cap \mathcal{P} \neq \emptyset) + \mathbb{P}(\mathcal{P}_n \cap \mathbf{B} \neq \mathcal{P} \cap \mathbf{B}) \\ &\leq e^{-L^2/36} + e^{-bn^{2\delta}} \leq e^{-L^2/36} + e^{-b(L/36)^2}, \end{aligned}$$

which yields the second part of Lemma 2.5.

□

Consider the random variable $m_r^3(Y^n(L))$ defined in (2.18). Assume we have a path $\gamma \in \mathcal{C}^r$. Then its animal $\mathbf{A}(\gamma)$ contains a subset $\mathbf{B} \in \Xi_{\mathbf{A}}^3$ with $m^3(Y^n(L), \mathbf{A})$ good boxes. By Lemma 2.1, if $\mathbf{B}_{\mathbf{z}}^{1/2,L}$ is a good box in $\mathbf{A}(\gamma)$ then γ has an edge lying in $\cup_{\bar{\mathbf{z}} \in \mathbf{C}_{\mathbf{z}}} \mathbf{B}_{\bar{\mathbf{z}}}^{1/2,L}$. On the other hand, if $\mathbf{B}_{\mathbf{z}_1}^{1/2,L}$ and $\mathbf{B}_{\mathbf{z}_2}^{1/2,L}$ are 3-distant then

$$\cup_{\bar{\mathbf{z}} \in \mathbf{C}_{\mathbf{z}_1}} \mathbf{B}_{\bar{\mathbf{z}}}^{1/2,L} \cap \cup_{\bar{\mathbf{z}} \in \mathbf{C}_{\mathbf{z}_2}} \mathbf{B}_{\bar{\mathbf{z}}}^{1/2,L} = \emptyset.$$

Thus, we must have that

$$m^3(Y^n(L), \mathbf{A}) \leq |\gamma| \leq r,$$

which implies that

$$\mathbb{P}(G_r^L(\mathcal{P}_n) > cr) \leq \mathbb{P}(m_{cr}^3(Y^n(L)) < r).$$

Together with Lemma 2.5 and Lemma (2.4) this proves (2.21) for large L_0 , which finishes the proof of Proposition 2.1. $\square$

Proof of Proposition 2.2. For this proof we say that $\mathbf{B}_{\mathbf{z}}^{1/2,L}$ is a good box, or equivalently $Z_{\mathbf{z}}^n(L) = 1$, if it is a full box. Then $Z^n(L) := \{Z_{\mathbf{z}}^n(L) : \mathbf{z} \in \mathbb{Z}^2\}$ is a collection of independent random variables.

Let $\mathbf{H}_j^n$ be the set of closed sites ($Z_{\mathbf{z}}^n(L) = 0$) in $\mathbb{Z}^2$ which are connected to $(j, 0)$, and

$$\bar{\mathbf{H}}_j^n := \{(0, j)\} \cup \mathbf{H}_j \cup \partial \mathbf{H}_j^n.$$

Also define $\mathbf{I}_s^n(Z^n(L)) := \cup_{j=1}^s \bar{\mathbf{H}}_j^n$.

By Lemma 2.1,

$$\gamma_r(\mathcal{P}_n) \subseteq \cup_{\mathbf{z} \in \mathbf{I}_{r/L}^n} \mathbf{B}_{\mathbf{z}}^{1/2,L},$$

and thus

$$|\gamma_r(\mathcal{P}_n)| \leq \sum_{\mathbf{z} \in \mathbf{I}_{r/L}^n} |\mathbf{B}_{\mathbf{z}}^{1/2,L} \cap \mathcal{P}_n|. \quad (2.24)$$

Since $\mathbb{P}(Z_{\mathbf{z}}^L(\mathcal{P}_n) = 0) = o(L)$ uniformly in n (by the same argument in Lemma 2.5), we can find L_0 sufficiently large so that $\mathbb{E}(\exp(|\mathbf{H}_0^n|)) < \infty$. This yields that

$$\mathbb{P}(|\mathbf{I}_s^n| > ys) \leq e^{c_1 y s}, \quad (2.25)$$

for $y > y_0$ and c_1, y_0 constants whose values do not depend on n .

From now on we fix such L_0 and write $M_s^n := M_s(Z^n(L_0))$ (defined in (2.15)) and $\mathbf{I}_s^n := \mathbf{I}_s^n(Z^n(L_0))$. By Lemma 2.3, for sufficiently small $c > 0$ one can find $b_1 > 0$ so that

$$\mathbb{P}(M_{cL_0 z r}^n \geq zr) \leq \mathbb{P}(M_{cL_0 z r}^\infty \geq (z-1)r) \leq e^{-b_1 z r}. \quad (2.26)$$

By (2.24),

$$\begin{aligned} \mathbb{P}(|\gamma_r(\mathcal{P}_n)| \geq zr) &\leq \\ \mathbb{P}(|\gamma_r(\mathcal{P}_n)| \geq zr, |\mathbf{I}_{r/L_0}^n| \leq cL_0 z(r/L_0)) &+ \mathbb{P}(|\mathbf{I}_{r/L_0}^n| > cL_0 z(r/L_0)) \leq \\ \mathbb{P}(M_{cL_0 z r}^n \geq zr) &+ \mathbb{P}(|\mathbf{I}_{r/L_0}^n| > cL_0 z(r/L_0)). \end{aligned}$$

Together with (2.26) and (2.25), this yields (2.22). □

Remark: the connective function on disordered graphs. Problems related to self-avoiding paths are connected with various branches of applied mathematics such as long chain polymers, percolation and ferromagnetism (Kesten [11], Hammersley [5]). One fundamental problem is the asymptotic behavior of the connective function κ_r defined by the logarithm of the number of self-avoiding paths (on some graph $\mathcal{G}$) starting at $\mathbf{v}$ and with r steps.

For planar and periodic graphs subadditivity arguments yields that $r^{-1}\kappa_r(\mathbf{v})$ converges, when $r \rightarrow \infty$, to some value $\kappa \in (0, \infty)$ (the connectivity constant) independent of the initial vertex $\mathbf{v}$. In disordered planar graphs subadditivity is lost but, if the underline graph possess some statistical symmetries (ergodicity), we may believed that the rescaled connective function still converges to some constant.

From Proposition 2.1 we obtain a linear upper bound for the connective function of the Delaunay triangulation, where we set $\mathbf{v}$ to be the closest vertex to the origin: there exist a constant $c > 0$ so that

$$\limsup_{r \rightarrow \infty} \frac{\kappa_r}{r} \leq c. \quad (2.27)$$

To prove (2.27) recall we have associate to each self-avoiding path γ and ordered sequence $\mathbf{A}(\gamma) = (\mathbf{z}_1, \dots, \mathbf{z}_n)$ of nearest neighbors integer sites, which maybe seen as animal in $\mathbb{Z}^2$ (Figure 3).

The number of self-avoiding paths associated to an ordered sequence $(\mathbf{z}_1, \dots, \mathbf{z}_n)$ should be at at most

$$|\mathbf{B}_{\mathbf{z}_1}^{1/2,1} \cap \mathcal{P}|! \times \dots \times |\mathbf{B}_{\mathbf{z}_n}^{1/2,1} \cap \mathcal{P}|!$$

($n!$ is factorial of n).

Now, every self-avoiding path with r steps intersects at most G_r boxes, and thus

$$N_r \leq 4^{G_r} \max_{|\mathbf{A}| \leq G_r} \{|\mathbf{B}_{\mathbf{z}_1}^{1/2,1} \cap \mathcal{P}|! \times \dots \times |\mathbf{B}_{\mathbf{z}_{|\mathbf{A}|}}^{1/2,1} \cap \mathcal{P}|!\},$$

which implies that

$$\kappa_r = \log N_r \leq (\log 4)G_r + \max_{|\mathbf{A}| \leq G_r} \left\{ \sum_{i=1}^{|\mathbf{A}|} \log |\mathbf{B}_{\mathbf{z}_i}^{1/2,1} \cap \mathcal{P}|! \right\}. \quad (2.28)$$

Denote

$$X_{\mathbf{z}} := \log |\mathbf{B}_{\mathbf{z}}^{1/2,1} \cap \mathcal{P}|!,$$

and let $X := \{X_{\mathbf{z}} : \mathbf{z} \in \mathbb{Z}^2\}$ and $M_s := M_s(X)$ (see (2.15)).

By (2.28), if $a > 1$ and $c < 1/2 \log 4$ then

$$\mathbb{P}(\kappa_r > ar) \leq \mathbb{P}(G_r > car) + \mathbb{P}(M_{car} > ar/2). \quad (2.29)$$

Since (2.16) holds for X , by Lemma 2.3 there exist $b_1, b_2 > 0$ such that if $a > 1$ and $c < \min\{1/2 \log 4, 1/2b_1\}$ then

$$\mathbb{P}(M_{car}^2 > ar/2) \leq e^{-b_2 car}. \quad (2.30)$$

By Proposition 2.1 we can find $c_3, c_4 > 0$ so that, if $ca > c_3$ then

$$\mathbb{P}(G_r > car) \leq e^{-c_4 r}.$$

Combining this with (2.29) and (2.30) we obtain that for $a > c_3 / \min\{1/2 \log 4, 1/2b_1\}$

$$\mathbb{P}(\kappa_r > ar) \leq e^{-b_2 car} + e^{-c_4 r}.$$

Together with Borel-Cantelli lemma, this proves (2.27).

3. DENSITY OF OPEN EDGES IN PERCOLATION AND THE TIME CONSTANT

In this section we prove a sufficient condition on the passage time distribution $\mathbb{F}$ to ensure that $\mu(\mathbb{F}) > 0$. We also show that the same condition ensures the existence of geodesics. These results will follow from the subsequent propositions.

Proposition 3.1. *There exist $p_0 \in (0, 1)$ so that if $p \in (p_0, 1)$ then*

$$\eta^*(p) = \liminf_{R \rightarrow \infty} \mathbb{P}_p^*(A_R) = 1.$$

In particular, $p_c^ < 1$.*

Let

$$t_r := \inf \left\{ \sum_{\mathbf{e} \in \gamma} \tau_{\mathbf{e}} : \gamma \in \mathcal{C}_r(\mathcal{P}_n) \right\}. \quad (3.31)$$

Proposition 3.2. *If $\mathbb{F}(0) < 1 - p_c^*$ then there exist constants $c_j > 0$ such that for all n sufficiently large, for all $r \geq 1$*

$$\mathbb{P}(t_r \leq c_1 r) \leq e^{-c_2 r}.$$

Proof of Proposition 3.1. Now we say that $\mathbf{B}_{\bar{\mathbf{z}}}^{1/2, L}$ is a good box, or equivalently $V_{\bar{\mathbf{z}}}(L, p) = 1$, if the following holds:

- $\mathbf{B}_{\bar{\mathbf{z}}}^{1/2, L}$ is a full box, with respect to $\mathcal{P}$, and $|\mathbf{B}_{\bar{\mathbf{z}}}^{1/2, L} \cap \mathcal{P}| \leq 4L^2$ for all $\bar{\mathbf{z}}$ with $|\bar{\mathbf{z}} - \mathbf{z}|_{\infty} \leq 1$;
- $X_{\mathbf{e}^*} = 1$ for all $\mathbf{e}^* \in \mathcal{V}_{\mathbf{e}}$ which intersects $\mathbf{B}_{\bar{\mathbf{z}}}^{1/2, L}$.

We denote $B_1(\mathbf{z}, L)$ and $B_2(\mathbf{z}, L)$ the event specified in the first and in the second item above, respectively. The law of large numbers implies that

$$\lim_{L \rightarrow \infty} \mathbb{P}(B_1(\mathbf{z}, L)) = 1. \quad (3.32)$$

Since $\mathbf{B}_{\bar{\mathbf{z}}}^{1/2,L}$ is a full box for all $\bar{\mathbf{z}}$ with $|\bar{\mathbf{z}} - \mathbf{z}|_\infty \leq 1$, by Lemma 2.1, the set of edges $\mathbf{e}^* \in \mathcal{V}_e$ intersecting $\mathbf{B}_{\bar{\mathbf{z}}}^{1/2,L}$ is contained in $\mathbf{B}_{\bar{\mathbf{z}}}^{3/2,L}$. Let V , E and F be the number of vertexes, edges and faces of the Delaunay triangulation on $\mathbf{B}_{\bar{\mathbf{z}}}^{3/2,L}$, respectively. Then

$$3E = 2F$$

and, by the Euler formula,

$$V - E + F = 2.$$

This yields that the number of edges in $\mathbf{B}_{\bar{\mathbf{z}}}^{3/2,L}$ has the same order as of vertexes in the same box. Thus, for some constant $b > 0$,

$$\mathbb{P}(B_1(\mathbf{z}, L) \cap B_2(\mathbf{z}, L)^c) \leq bL^2(1 - p).$$

Together with (3.32), this yields that for all $\delta > 0$ we can chose L sufficiently large, and then $p = p(L, \delta)$ sufficiently close to 1, to have that

$$\mathbb{P}(\mathbf{B}_{\mathbf{z}}^{1/2,L} \text{ is a good box}) > 1 - \delta. \quad (3.33)$$

From Lemma 2.2 we have that $V := \{V_{\mathbf{z}}(L, p) : \mathbf{z} \in \mathbb{Z}^2\}$ is a collection of 3-dependent random variables.

Notice that by a similar argument to construct γ_r (see subsection 2.3) and by using Lemma 2.1 one can get that

$$\begin{aligned} & [\exists \text{ a crossing composed by good boxes of } [0, 3R/L] \times [0, R/L]] \subseteq \\ & [\exists \text{ an open crossing, with respect to } X, \text{ of } [0, 3R] \times [0, R]]. \end{aligned}$$

Combining (3.33) with 3-dependence, one can find L_0 large enough and then p_0 sufficiently close to 1 so that the probability of the first event in the above inclusion goes to 1 as R goes to infinity, which finishes the proof of Proposition 3.1. $\square$

Proof of Proposition 3.2. First we claim it suffices to prove Proposition 3.2 if $\mathbb{P}(\tau_{\mathbf{e}} = 1) = p = 1 - \mathbb{P}(\tau_{\mathbf{e}} = 0)$.

To see this, assume that $\mathbb{F}(0) < 1 - p_c^*$. Since $\mathbb{F}$ is right-continuous, we can chose $\epsilon > 0$ small so that $\mathbb{F}(\epsilon) < 1 - p_c^*$. Let

$$\tau_e^\epsilon := \mathbb{I}(\tau_{\geq \epsilon})$$

and denote t_r^ϵ the infimum defined in (3.31) but considering the travel times τ_e^ϵ . Therefore

$$\mathbb{P}(\tau_e^\epsilon = 0) < 1 - p_c^* \text{ and } \epsilon t_r^\epsilon \leq t_r,$$

which yields the last claim.

Now assume that $\mathbb{P}(\tau_{\mathbf{e}} = 1) = p = 1 - \mathbb{P}(\tau_{\mathbf{e}} = 0)$. Let $L > 0$ and $\mathbf{z} \in \mathbb{Z}^2$ and define that $\mathbf{B}_{\mathbf{z}}^{1/2,L}$ is a good box if the following holds:

- For all $\mathbf{z}' \in \mathbb{Z}^2$ with $|\mathbf{z} - \mathbf{z}'| = 2$, $\mathbf{B}_{\mathbf{z}'}^{1/2,L}$ is a full box;

- For all $\gamma \in \mathcal{C}(\mathbf{B}_{\mathbf{z}}^{1/2,L}, \partial\mathbf{B}_{\mathbf{z}}^{3/2,L})$ we have $t(\gamma) \geq 1$.

Let

$$W_{\mathbf{z}}^n(L) := \mathbb{I}(\mathbf{B}_{\mathbf{z}}^{1/2,L} \text{ is a good box}),$$

and

$$p(n, L) := \inf_{\mathbf{z} \in \mathbb{Z}^2} \mathbb{P}(W_{\mathbf{z}}^n(L) = 1).$$

Lemma 3.1. $W_{\mathbf{z}}^n(L) := \{W_{\mathbf{z}}^n(L) : \mathbf{z} \in \mathbb{Z}^2\}$ is 5-dependent for all $L > 0$. Moreover, if $\mathbb{P}(\tau_{\mathbf{e}} = 0) < 1 - p_c^*$ then for all $\epsilon > 0$ there exists $L_{\epsilon} > 0$ (independent of n) so that for all $n \geq L_{\epsilon}$

$$1 - \epsilon < p(n, L_{\epsilon}) \leq 1.$$

Proof of Lemma 3.1. 5-dependence follows from Lemma 2.2.

Now consider the event

$$H_{\mathbf{z}}^{L,n} := [\mathbf{B}_{\mathbf{z}'}^{1/2,L} \text{ is a full box (with respect to } \mathcal{P}_n) \forall |\mathbf{z} - \mathbf{z}'| = 2]$$

and the event

$$G_{\mathbf{z}}^{L,n} := [t(\gamma) \geq 1 \forall \gamma \in \mathcal{C}(\mathbf{B}_{\mathbf{z}}^{1/2,L}, \partial\mathbf{B}_{\mathbf{z}}^{3/2,L})].$$

Then

$$\mathbb{P}(W_{\mathbf{z}}^n(L) = 0) \leq \mathbb{P}((H_{\mathbf{z}}^{L,n})^c) + \mathbb{P}((G_{\mathbf{z}}^{L,n})^c). \quad (3.34)$$

As we have seen before,

$$\mathbb{P}((H_{\mathbf{z}}^{L,n})^c) \rightarrow 0 \text{ when } L \rightarrow \infty \text{ uniformly in } n.$$

This yields Lemma 3.1 if we prove that if $\mathbb{P}(\tau_{\mathbf{e}} = 0) < 1 - p_c^*$ then for all $\epsilon > 0$ there exists $L_{\epsilon} > 0$ so that for all $n \geq L_{\epsilon}$

$$1 - \epsilon < \mathbb{P}(G_{\mathbf{z}}^{L_{\epsilon},n}) \leq 1. \quad (3.35)$$

We first prove (3.35) for $n = \infty$. Let $X_{\mathbf{e}^*} := \tau_{\mathbf{e}}$, where $\mathbf{e}^*$ is the edge in $\mathcal{V}_{\mathbf{e}}$ (with point configuration $\mathcal{P}$) dual to $\mathbf{e}$. Then $\{X_{\mathbf{e}^*} : \mathbf{e}^* \in \mathcal{V}_{\mathbf{e}}\}$ defines a bond percolation model on $\mathcal{V}$ with law $\mathbb{P}_p^*$.

Let

$$R_L^1 := [L/2, 3L/2] \times [-3L/2, 3L/2], \quad R_L^2 := [-3L/2, 3L/2] \times [L/2, 3L/2],$$

$$R_L^3 := [-3L/2, -L/2] \times [-3L/2, 3L/2], \quad R_L^4 := [-3L/2, 3L/2] \times [-3L/2, -L/2].$$

Let A_L^i be the event that there exists an open crossing in $\mathcal{V}$, as in the definition of the event A_L , but now translate to the rectangle R_L^i .

Denote by F_L the event that an open circuit σ^* in $\mathcal{V}$ which surrounds $\mathbf{B}_{\mathbf{0}}^{1/2,L}$ and lies inside $\mathbf{B}_{\mathbf{0}}^{3/2,L}$ does not exist. Thus one can see that

$$\cap_{i=1}^4 A_L^i \subseteq (F_L)^c.$$

Notice also that, if there exists an open circuit σ^* in $\mathcal{V}$ which surrounds $\mathbf{B}_0^{1/2,L}$ and lies inside $\mathbf{B}_0^{3/2,L}$ then every path γ connecting $\mathbf{B}_0^{1/2,L}$ to $\partial\mathbf{B}_0^{3/2,L}$ has an edge crossing with σ^* , and thus $t(\gamma) \geq 1$. Together with translation invariance, this yields

$$\mathbb{P}((G_0^{L,\infty})^c) \leq \mathbb{P}_p^*(F_L) \leq 4(1 - \mathbb{P}_p^*(A_L)). \quad (3.36)$$

Since $p > p_c^*$, by using (3.36) and the definition of p_c^* one can obtain (3.35) for $n = \infty$.

Now,

$$\mathbb{P}(\mathcal{P}_\infty \neq \mathcal{P}_n \text{ in } \mathbf{B}_{\mathbf{z}}^{5/2,L}) \leq c_1 L^2 e^{-c_2 n^{2\delta}}.$$

for some constant $c_1 > 0$. Thus, for all $n \geq L$

$$\mathbb{P}((G_{\mathbf{z}}^{L,n})^c) \leq \mathbb{P}((G_{\mathbf{z}}^{L,\infty})^c) + c_1 L^2 e^{-c_2 L^{2\delta}}. \quad (3.37)$$

By (3.36) and (3.37), given $\epsilon > 0$ we can find L_ϵ so that for all $n \geq L_\epsilon$

$$\mathbb{P}((G_{\mathbf{z}}^{L_\epsilon,n})^c) \leq \epsilon,$$

which finishes the proof of this lemma. □

Now consider the random variable $m_r^5(W^n(L))$ defined by (2.18). Combining Lemma 2.4 with Lemma 3.1, one obtains that there exists $L_0 > 0$ and $b_1, b_2 > 0$, whose value depends only on L_0 , such that for all $n \in [1, \infty]$, for all $r > 0$

$$\mathbb{P}(m_r^5(W^n(L_0)) \leq b_1 r) \leq e^{-b_2 r}. \quad (3.38)$$

Let $\gamma \in \mathcal{C}_r$ and consider the animal $\mathbf{A}(\gamma)$ previously defined. Then $\mathbf{A}(\gamma)$ contains a subset $\mathbf{B} \in \Xi_{\mathbf{A}}^5$ with $m^5(\mathbf{A}(\gamma), W^n(L))$ good boxes. If $\mathbf{B}_{\mathbf{z}_1}^{1/2,L}$ and $\mathbf{B}_{\mathbf{z}_2}^{1/2,L}$ are 5-distant good boxes in $\mathbf{A}(\gamma)$ then (by definition) for each $j = 1, 2$ there exist a piece of γ , say γ_j , connecting $\partial\mathbf{B}_{\mathbf{z}_j}^{1/2,L}$ to $\partial\mathbf{B}_{\mathbf{z}_j}^{3/2,L}$ and so that $t(\gamma_j) \geq 1$. Since these good boxes are 5-distant, by Lemma 2.1, γ_1 and γ_2 must be disjoint. This yields that $t(\gamma) \geq 2$. By repeating this argument inductively, one gets that

$$m^5(\mathbf{A}(\gamma), W^n(L_0)) \leq t(\gamma) \leq r.$$

Therefore,

$$\mathbb{P}(t_r \leq cr) \leq \mathbb{P}(g_r < br) + \mathbb{P}(m_{br}^5(W^n(L)) \leq cr).$$

Together with (3.38) and Proposition 2.1, this yields Proposition 3.2. □

Corollary 3.1. *If $\mathbb{F}(0) < 1 - p_c^*$ then there exist constants $c_j > 0$ such that for all n sufficiently large, for all $r > 0$*

$$\mathbb{P}(T(\mathbf{0}, \mathbf{r}) < c_1 r) \leq e^{-c_2 r}. \quad (3.39)$$

In particular, if $\mathbb{F}(0) < 1 - p_c^$ then $\mu(\mathbb{F}) > 0$.*

Proof of Corollary 3.1. If we have a path γ connecting $\mathbf{0}$ to $\mathbf{r}$ then it intersects at least r unit boxes $\mathbf{B}_{\mathbf{z}}^{1/2,1}$. Thus, if $b > 0$ then

$$\mathbb{P}(T(\mathbf{0}, \mathbf{r}) < c_1 r) \leq \mathbb{P}(G_{br} > r) + \mathbb{P}(t_{br} \leq c_1 r).$$

Combining Proposition 3.2 with Proposition 2.1, one can get (3.1).

The second part follows from (3.1), together with Borel-Cantelli Lemma.

□

Corollary 3.2. *If $\mathbf{F}(0) < 1 - p_c^*$ then for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^2$ $\mathbb{P}$ -a.s. there exist at least one geodesic connecting $\mathbf{x}$ to $\mathbf{y}$. Further, if $\mathbb{F}$ is continuous, then we have $\mathbb{P}$ -a.s. uniqueness.*

Proof of Corollary 3.2. Let

$$d_r(\mathbf{v}) := \inf\{T(\mathbf{v}, \bar{\mathbf{v}}) : |\bar{\mathbf{v}}| \geq r\}.$$

By Proposition 2.1 and Proposition 3.2, we can find sufficiently small $c > 0$ so that $\mathbb{P}$ -a.s.

$$0 < c < \liminf_r \frac{d_r(\mathbf{v})}{r}.$$

In particular, $d_r(\mathbf{v}) \rightarrow \infty$ when $r \rightarrow \infty$.

Therefore, if we fix $\mathbf{v}, \bar{\mathbf{v}} \in \mathcal{P}_n$ then we can find r large enough so that $d_r(\mathbf{v}) > T(\mathbf{v}, \bar{\mathbf{v}}) + 1$. This implies that the passage time between $\mathbf{v}$ and $\bar{\mathbf{v}}$ will be attained by a path in the finite collection of paths connecting $\mathbf{v}$ to $\bar{\mathbf{v}}$ and lying inside the ball centered at $\mathbf{v}$ and with radius r .

To complete the proof, notice that if $\mathbb{F}$ is a continuous function then $\mathbb{P}$ -a.s. there are no finite path γ and $\bar{\gamma}$ with $\sum_{\mathbf{e} \in \gamma} \tau_{\mathbf{e}} = \sum_{\mathbf{e} \in \bar{\gamma}} \tau_{\mathbf{e}}$.

□

4. MARTINGALES WITH BOUNDED INCREMENTS AND THE RATE OF CONVERGENCE

We formulate the abstract martingale estimate used by Howard and Newman [8].

Lemma 4.1. *Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and let $\{\mathcal{F}_k\}_{k \geq 0}$ be an increasing family of σ -algebras of measurable sets. Let $\{M_k\}_{k \geq 0}$, $M_0 = 0$, be a martingale with respect to the filtration $\{\mathcal{F}_k\}_{k \geq 0}$ and let $\{U_k\}_{k \geq 0}$ be a collection of positive random variables that are $\mathcal{F}$ -measurable. Assume that the increments $\Delta_k = M_k - M_{k-1}$ satisfy*

$$|\Delta_k| \leq c, \text{ for some constant } c \geq 0, \tag{4.40}$$

and

$$E(\Delta_k^2 \mid \mathcal{F}_{k-1}) \leq E(U_k \mid \mathcal{F}_{k-1}). \tag{4.41}$$

Assume further that for some finite constants $0 < c_1, v, x_0$ with $x_0 \geq c^2$, we have that for all $x \geq x_0$,

$$P\left(\sum_{k \geq 1} U_k > x\right) \leq c_1 \exp(x^v). \quad (4.42)$$

Then almost surely $M = \lim_{k \rightarrow \infty} M_k$ exists. Moreover, there exists constants $0 < c_2, c_3$, whose values do not depend on c and x_0 , such that for all $x \leq x_0^v$,

$$P(|M| > x\sqrt{x_0}) \leq c_2 \exp(-c_3 x). \quad (4.43)$$

Proof of Lemma 4.1. See the proof of Lemma 5.2 in Howard and Newman [8]

□

We construct the underline probability space as follows. For each $k \geq 1$ let

$$I_k := \{(a, b, c) : a \geq 1, b \geq k \text{ and } c \geq 1 \text{ if } b \neq k \text{ or } c > a \text{ if } b = k\},$$

and denote

$$\omega_k := (N_k, (U_{k,a})_{a \geq 1}, (\tau_{k,a}^{b,c})_{(a,b,c) \in I_k})$$

where N_k is a Poisson random variable with intensity $n^{2\delta}$, $(U_{k,a})_{a \geq 1}$ is a sequence of independent points uniformly distributed in $\mathbf{B}_k^n = \mathbf{B}_{\mathbf{u}_k}^{1/2, n^\delta}$, and $(\tau_{k,a}^{b,c})_{(a,b,c) \in I_k}$ is a sequence of independent random variables with distribution $\mathbb{F}$.

Let $(\Omega_k, \mathbb{P}_k, \mathcal{A}_k)$ be the probability space induced by ω_k and set $\omega = (\omega_k)_{k \geq 1}$ and

$$\mathbb{P} = \prod_{k=1}^{\infty} \mathbb{P}_k.$$

To determine the point process $\mathcal{P}$, we put N_k points in the box $\mathbf{B}_k^n$ given by $U_{k,1}, \dots, U_{k,N_k}$. For each $\mathbf{e} \in \mathcal{D}_e$ we know there exists $k \geq 1$ and $(a, b, c) \in I_k$ so that $\mathbf{e} = (U_{k,a}, U_{b,c})$ and so we set $\tau_{\mathbf{e}} := \tau_{k,a}^{b,c}$.

We shall use the following σ -fields:

$$\mathcal{F}_0 := \{\emptyset, \Omega\}$$

$$\mathcal{F}_k := \sigma\text{-field generated by } \omega_1, \dots, \omega_k.$$

Let $T_n := T(\mathbf{0}, \mathbf{n})$. The martingale representation of $T_n - \mathbb{E}T_n$ is

$$T_n - \mathbb{E}T_n = \sum_{k=1}^{\infty} \{\mathbb{E}(T_n | \mathcal{F}_k) - \mathbb{E}(T_n | \mathcal{F}_{k-1})\}.$$

This representation is valid because $M_0 := 0$ and

$$\begin{aligned} M_l &:= \sum_{k=1}^l \{\mathbb{E}(T_n | \mathcal{F}_k) - \mathbb{E}(T_n | \mathcal{F}_{k-1})\} \\ &= \mathbb{E}(T_n | \mathcal{F}_l) - \mathbb{E}T_n \end{aligned}$$

defines a $\mathcal{F}_k$ -martingale that, under (1.5), converges a.s. (and also in L^2), to $T_n - \mathbb{E}T_n$.

The increments of M_l are

$$\Delta_k = \mathbb{E}(T_n \mid \mathcal{F}_k) - \mathbb{E}(T_n \mid \mathcal{F}_{k-1}),$$

and the main step is to estimate $\mathbb{E}(\Delta_k^2 \mid \mathcal{F}_{k-1})$.

4.1. Successive approximations. In order to satisfies the prescription of Lemma 4.1 we shall modified the configuration ω . We define

$$\bar{\omega}_k := \bar{\omega}_k(n, \omega_k)$$

as follows: we change the point configuration $\mathcal{P} \cap \mathbf{B}_k^n$ to $\mathcal{P}_n \cap \mathbf{B}_k^n$; the travel time configuration is changed by putting travel times ³

$$\bar{\tau}_{k,a}^{b,c} := \min\{\tau_{k,a}^{b,c}, 8b^{-1} \log n\}.$$

We note that Proposition 3.2 still holds for $t_r(\bar{\mathcal{P}}, (\bar{\tau}_e)_{\bar{\mathcal{D}}_e})$ (with constants not depending on n), since for large enough n , $\bar{\tau}_e \geq \min\{\tau_e, 1\}$.

We denote $\bar{\omega} := (\bar{\omega}_k)_{k \geq 1}$. When a variable, such as travel times, geodesics and increments, is a function of $\bar{\omega}$ we will decorate it with a bar. For instance we write $\bar{T}_r(\omega) := T_r(\bar{\omega})$ and $\bar{\rho}_r := \rho_r(\bar{\omega})$.

Lemma 4.2. *If $\mathbb{F}(0) < 1 - p_c^*$ and (1.5) holds then,*

$$\mathbb{P}(|T_n - \bar{T}_n| > x) \leq e^{-c_1 n^{2\delta}} + e^{-c_2 x} \quad (4.44)$$

and

$$\mathbb{E}(\{T_n - \bar{T}_n\}^2) \leq c_3. \quad (4.45)$$

Proof of Lemma 4.2. To prove (4.44) we first claim that

$$\mathbb{P}(\bar{\rho}_n \not\subseteq [-cn, cn]^2) \leq e^{-c_2 n} \quad (4.46)$$

(for sufficiently large $c > 0$).

Lemma 4.3. *Under (1.5), there exists $x_0, c_0 > 0$ so that for all $x > x_0$*

$$\mathbb{P}(\bar{T}_r > xr) \leq e^{-c_0 xr}.$$

Proof of Lemma 4.3. By Proposition 2.2, there exist $c_1, z_0 > 0$ so that if $cx > z_0$ then

$$\mathbb{P}(\bar{T}_r > xr) \leq \mathbb{P}(\bar{T}_r > xr, |\bar{\gamma}_r| \leq cxr) + e^{-c_1 cxr}. \quad (4.47)$$

By independence,

$$\mathbb{P}(\bar{T}_r > xr, |\bar{\gamma}_r| \leq cxr) \leq \sum_{j=0}^{cxr} \mathbb{P}\left(\sum_{\mathbf{e} \in \bar{\gamma}_r} \bar{\tau}_e > xr \mid |\bar{\gamma}_r| = j\right) \mathbb{P}(|\bar{\gamma}| = j) \leq$$

³ $b > 0$ is given by (1.5)

$$\sum_{j=0}^{cxr} \mathbb{P}\left(\sum_{i=0}^j \bar{\tau}_i > xr\right) \mathbb{P}(|\bar{\gamma}| = j) \leq e^{-xar} (\mathbb{E}e^{a\tau})^{cx} \leq e^{-axr/2}$$

if $c < a \log \mathbb{E}e^{a\tau}$. Together with 4.47, this yields Lemma 4.3.

□

To prove (4.46) notice that,

$$\begin{aligned} \mathbb{P}(\bar{\rho}_n \not\subseteq [-cn, cn]^2) &\leq \\ \mathbb{P}(|\bar{\rho}_n| > bn) + \mathbb{P}(\exists \gamma \in \mathcal{C}^{bn} : \gamma \not\subseteq [-cn, cn]^2). \end{aligned} \quad (4.48)$$

Now,

$$\begin{aligned} \mathbb{P}(|\bar{\rho}_n| > bn) &\leq \mathbb{P}(\bar{T}_n > an) + \mathbb{P}(\bar{T}_n \leq an \text{ and } |\bar{\rho}_n| > bn) \\ &= \mathbb{P}(\bar{T}_n > an) + \mathbb{P}(t_{bn} \leq \frac{a}{b}bn). \end{aligned}$$

Together with Proposition 3.2 and Lemma 4.3, this yield that if we take $a > x_0$, $b > 0$ so that $a/b < c_1$ (given by Proposition 3.2) then

$$\mathbb{P}(|\bar{\rho}_n| > bn) \leq e^{c_2n}, \quad (4.49)$$

for some constant $c_2 > 0$.

By Proposition 2.1, if we take c sufficiently large,

$$\mathbb{P}(\exists \gamma \in \mathcal{C}^{bn} : \gamma \not\subseteq [-cn, cn]^2) \leq \mathbb{P}(G_{bn} \geq cn)$$

will also decays exponentially fast with n . Together with (4.48) and (4.49), this proves (4.46).

Turning back to the proof of (4.44), assume that $\bar{\rho}_n \subseteq [-cn, cn]^2$ and $\mathcal{P} = \mathcal{P}_n$. Thus

$$|\bar{T}_n - T_n| \leq |\bar{T}_n| + |T_n| \leq 2t(\bar{\gamma})_n.$$

This yields,

$$\begin{aligned} \mathbb{P}(|\bar{T}_n - T_n| > x) &\leq \mathbb{P}(\mathcal{P} \neq \mathcal{P}_n \text{ in } [-c_1n, c_1n]^2) + \\ &\mathbb{P}(\bar{\rho}_n \not\subseteq [-cn, cn]^2) + \mathbb{P}\left(\sum_{[-c_1n, c_1n]^2} \tau_e \mathbf{I}(\tau_e > 8a^{-1} \log n) > x\right). \end{aligned} \quad (4.50)$$

Notice that, since the mean number of Poisson points inside B_k^n is $n^{2\delta}$, by using simple large deviations results for Poisson process we have that for each $b > 0$ there exists a constant $b_1 = b_1(b) > 0$ such that for all $n \in [1, \infty)$

$$\mathbb{P}(\mathcal{P} \cap [-bn, bn]^2 \neq \mathcal{P}_n \cap [-bn, bn]^2) \leq e^{-b_1n^{2\delta}}. \quad (4.51)$$

Let $\mathbf{B} \subseteq \mathbb{R}^2$ and denote $E_{\mathbf{B}}(\mathcal{P})$ the number of edges $(\mathbf{v}, \mathbf{v}')$ in $\mathcal{D}$ so that $\mathbf{v}, \mathbf{v}' \in \mathbf{B}$. Since the number of edges is comparable with the number of vertexes (see the proof of

Proposition 3.1) we have that for each $b > 0$ there exists $b_1 = b_1(b) > 0$ such that for all $r > 0$

$$\mathbb{P}(E_{[-br, br]^2}(\mathcal{P}) > 2(br)^2) \leq e^{-b_1 r^2}. \quad (4.52)$$

To estimate the last term in the right hand side of (4.50) notice that, by (4.52),

$$\begin{aligned} & \mathbb{P}\left(\sum_{[-c_1 n, c_1 n]^2} \tau_e \mathbf{I}(\tau_e > 8a^{-1} \log n) > x\right) \leq \\ & \mathbb{P}\left(\sum_{[-c_1 n, c_1 n]^2} \tau_e \mathbf{I}(\tau_e > 8a^{-1} \log n) > x \mid E_{[-c_1 n, c_1 n]^2} \leq b_1 n^2\right) + e^{b_2 n^2}. \end{aligned}$$

By using a similar procedure presented in Kesten [13] (see (2.37) there) one can prove that

$$\mathbb{P}\left(\sum_{[-c_1 n, c_1 n]^2} \tau_e \mathbf{I}(\tau_e > 8a^{-1} \log n) > x \mid E_{[-c_1 n, c_1 n]^2} \leq b_1 n^2\right) \leq e^{-c_1 x}.$$

Together with (4.46), (4.50) and (4.51), this yields (4.44).

By Lemma 4.3, we also have that

$$\mathbb{P}(|\bar{T}_n - T_n| > ny) \leq \mathbb{P}(\bar{T}_n > ny/2) + \mathbb{P}(T_n > ny/2) \leq e^{-by_n}, \quad (4.53)$$

if $y \geq y_0$.

Thus

$$\begin{aligned} \mathbb{E}(\{\bar{T}_n - T_n\}^2) &= 2n^2 \int_0^\infty \mathbb{P}(|\bar{T}_n - T_n| > ny) y dy \\ &= 2n^2 \left[\int_0^{y_0} \mathbb{P}(|\bar{T}_n - T_n| > ny) y dy + \int_{y_0}^\infty \mathbb{P}(|\bar{T}_n - T_n| > ny) y dy \right]. \end{aligned}$$

By (4.44) and (4.50), the last term is bounded by a constant and this completes the proof of Lemma 4.2

□

4.2. Main estimate. We introduce the the following notation: if $\omega = (\omega_k), \sigma = (\sigma_k) \in \Omega$ then we denote

$$[\omega, \sigma]_k := (\omega_1, \dots, \omega_k, \sigma_{k+1}, \sigma_{k+2}, \dots).$$

We also set

$$\nu_k := \prod_{j=k}^\infty \mathbb{P}_k.$$

Since

$$\int \bar{T}_n[\omega, \sigma]_k \nu_{k+1}(d\sigma) = \int \bar{T}_n[\omega, \sigma]_k \nu_k(d\sigma),$$

we have

$$\bar{\Delta}_k = \int \bar{T}_n[\omega, \sigma]_k \nu_{k+1}(d\sigma) - \int \bar{T}_n[\omega, \sigma]_{k-1} \nu_k(d\sigma)$$

$$= \int \{\bar{T}_n[\omega, \sigma]_k - \bar{T}_n[\omega, \sigma]_{k-1}\} \nu_k(d\sigma).$$

Let F_k be the event that a geodesic from $\mathbf{0}$ to $\mathbf{n}$ has a vertex in $\mathbf{B}_k^n$. Let I_k be the indicator function of F_k and denote $\bar{I}_k(\omega) := I_k(\bar{\omega})$.

Lemma 4.4.

$$|\bar{T}_n[\omega, \sigma]_k - \bar{T}_n[\omega, \sigma]_{k-1}| \leq c_1 n^{2\delta} \log n \times \max\{\bar{I}_k[\omega, \sigma]_k, \bar{I}_k[\omega, \sigma]_{k-1}\}.$$

Proof of Lemma 4.4. By Lemma 2.1, if $\bar{I}_k[\omega, \sigma]_k = 0$ then $\bar{\rho}_n[\omega, \sigma]_k$ is also a path in $\mathcal{D}[\omega, \sigma]_{k-1}$. Thus

$$T_n[\omega, \sigma]_{k-1} \leq t(\rho_n[\omega, \sigma]_k) = T_n[\omega, \sigma]_k.$$

Analogously, if $\bar{I}_k[\omega, \sigma]_{k-1} = 0$ then

$$T_n[\omega, \sigma]_k \leq t(\rho_n[\omega, \sigma]_{k-1}) = T_n[\omega, \sigma]_{k-1}.$$

Therefore,

$$\max\{\bar{I}_k[\omega, \sigma]_k, \bar{I}_k[\omega, \sigma]_{k-1}\} = 0 \Rightarrow |\bar{T}_n[\omega, \sigma]_k - \bar{T}_n[\omega, \sigma]_{k-1}| = 0. \quad (4.54)$$

Now assume $\bar{I}_k[\omega, \sigma]_k = 1$. Let $\mathbf{v}_f$ (resp. $\mathbf{v}_l$) be the first (resp. the last) vertex $\mathbf{v} \in \rho_n[\omega, \sigma]_k$ so that $\mathbf{C}_{\mathbf{v}} \cap \mathbf{B}_{\mathbf{z}_k}^{13, n^{2\delta}}$. Define

$$\psi_n := \rho(\mathbf{0}, \mathbf{v}_f) \gamma(\mathbf{v}_f, \mathbf{v}_l) \rho(\mathbf{v}_l, \mathbf{n})$$

which is the concatenation of these three paths.

By Lemma 2.1 and the definition of $\mathcal{P}_n$, which ensures that $\mathbf{B}_{\mathbf{u}_k}^{13, n^{2\delta}}$ has enough points to isolate $\mathbf{B}_k^n$ from $\mathcal{G}_{\mathcal{P}_n}((\mathbf{B}_{\mathbf{u}_k}^{13, n^{2\delta}})^c)$ (see sub-section 2.1), $\mathcal{D}[\omega, \sigma]_k$ and $\mathcal{D}[\omega, \sigma]_{k-1}$ coincide outside $\mathbf{B}_{\mathbf{u}_k}^{13, n^{2\delta}}$. Thus, ψ_n is also a path in $\mathcal{D}[\omega, \sigma]_{k-1}$ and so

$$\begin{aligned} T_n[\omega, \sigma]_{k-1} &\leq t(\psi_n) = T(\mathbf{0}, \mathbf{v}_f)[\omega, \sigma]_k + t(\gamma(\mathbf{v}_f, \mathbf{v}_l))[\omega, \sigma]_{k-1} + T(\mathbf{v}_l, \mathbf{n})[\omega, \sigma]_k \\ &\leq T_n[\omega, \sigma]_k + t(\gamma(\mathbf{v}_f, \mathbf{v}_l))[\omega, \sigma]_{k-1}. \end{aligned}$$

Analogously, if $\bar{I}_k[\omega, \sigma]_{k-1} = 1$, then

$$\begin{aligned} T_n[\omega, \sigma]_k &\leq t(\psi_n) = T(\mathbf{0}, \mathbf{v}_f)[\omega, \sigma]_{k-1} + t(\gamma(\mathbf{v}_f, \mathbf{v}_l))[\omega, \sigma]_k + T(\mathbf{v}_l, \mathbf{n})[\omega, \sigma]_{k-1} \\ &\leq T_n[\omega, \sigma]_{k-1} + t(\gamma(\mathbf{v}_f, \mathbf{v}_l))[\omega, \sigma]_k. \end{aligned}$$

Therefore,

$$|\bar{T}_n[\omega, \sigma]_k - \bar{T}_n[\omega, \sigma]_{k-1}| \leq$$

$$\max\{t(\gamma(\mathbf{v}_f, \mathbf{v}_l))[\omega, \sigma]_k, t(\gamma(\mathbf{v}_f, \mathbf{v}_l))[\omega, \sigma]_{k-1}\} \max\{\bar{I}_k[\omega, \sigma]_{k-1}, \bar{I}_k[\omega, \sigma]_k\}. \quad (4.55)$$

Lemma 2.1 and the definition of $\mathcal{P}_n$ also ensures that,

$$\max\{|t(\gamma(\mathbf{v}_f, \mathbf{v}_l))[\omega, \sigma]_k|, |t(\gamma(\mathbf{v}_f, \mathbf{v}_l))[\omega, \sigma]_{k-1}|\} \leq c_1 n^{2\delta},$$

for some constant $c_1 > 0$. Together with

$$\bar{\tau}_e \leq 8a^{-1} \log n$$

and (4.55), this implies Lemma 4.4. □

Let $U_k := 2(c_1)^2 n^{4\delta} (\log n)^2 \bar{I}_k$, where c_1 is the constant given by Lemma 4.4.

Lemma 4.5. *If $\mathbb{F}(0) < 1 - p_c^*$ and (1.5) holds then for all sufficiently large n*

$$\mathbb{E}\left(\sum_{k=1}^{\infty} U_k\right) \leq n^{1+5\delta}. \quad (4.56)$$

Further, for all $\delta \in (0, 1/8)$ for all $y \geq n^{1+7\delta}$,

$$\mathbb{P}\left(\sum_{k=1}^{\infty} U_k > y\right) \leq e^{-c_2 y^{1/2}}. \quad (4.57)$$

Proof of Lemma 4.5. By definition

$$\sum_{k=1}^{\infty} U_k = c_1 n^{4\delta} (\log n)^2 \sum_{k=1}^{\infty} I_k \leq c_1 n^{4\delta} (\log n)^2 \sup |\bar{\rho}_n|. \quad (4.58)$$

Thus

$$\begin{aligned} \mathbb{P}\left(\sum_{k=1}^{\infty} I_k > ny\right) &\leq \mathbb{P}(\sup |\bar{\rho}_n| > ny) \\ &\leq \mathbb{P}(\bar{T}_n \leq cny) + \mathbb{P}(t_{ny} < cny). \end{aligned}$$

By Proposition 3.2, if we chose $c > 0$ sufficiently small,

$$\mathbb{P}(t_{ny} < cny) \leq e^{\bar{c} y^n}.$$

Therefore,

$$\begin{aligned} \mathbb{E}\left(\sum_{k=1}^{\infty} I_k\right) &\leq \mathbb{E}(\sup |\bar{\rho}_n|) = n \int_0^{\infty} \mathbb{P}(\sup |\bar{\rho}_n| > ny) dy \\ &\leq n \int_0^{\infty} \mathbb{P}(\bar{T}_n \geq cny) dy + \int_0^{\infty} \mathbb{P}(t_{ny} < cny) dy \\ &= \mathbb{E}(\bar{T}_n) + \tilde{c}n, \end{aligned}$$

which proves (4.56).

From Proposition 3.2 and Lemma 4.3, if $x > c_0 n$ then

$$\mathbb{P}(\sup |\bar{\rho}_n| > x) \leq \mathbb{P}(\bar{T}_n \geq cx) + \mathbb{P}(t_x < cx) \leq e^{bx}.$$

Together with (4.56), this yields (4.57).

□

Proof of Theorem 1. (1.4) follows from Corollary 3.1.

Now we prove (1.6): by Lemma 4.4, the Schwarz inequality and the Fubini theorem,

$$\begin{aligned} \mathbb{E}(\bar{\Delta}_k^2 \mid \mathcal{F}_{k-1}) &= \int \left\{ \int \{\bar{T}_n[\omega, \sigma]_k - \bar{T}_n[\omega, \sigma]_{k-1}\} \nu_k(d\sigma) \right\}^2 \mathbb{P}_k(d\omega) \\ &\leq c_1^2 n^{4\delta} (\log n)^2 \int \left\{ \int \max\{\bar{I}_k[\omega, \sigma]_k, \bar{I}_k[\omega, \sigma]_{k-1}\} \nu_k(d\sigma) \right\} \mathbb{P}_k(d\omega) \\ &\leq 2c_1^2 n^{4\delta} (\log n)^2 \int \bar{I}_k[\omega, \sigma]_{k-1} \nu_k(d\sigma) = \mathbb{E}(U_k \mid \mathcal{F}_{k-1}). \end{aligned} \quad (4.59)$$

From Lemma 4.5 and (4.59), for all n sufficiently large

$$\mathbb{V}\bar{T}_n = \sum_{k=1}^{\infty} \mathbb{E}(\bar{\Delta}_k^2) \leq \mathbb{E}\left(\sum_{k=1}^{\infty} U_k\right) \leq n^{1+5\delta}. \quad (4.60)$$

Now, Lemma 4.2 and the Schwarz inequality implies that

$$\begin{aligned} \mathbb{V}T_n - \mathbb{V}\bar{T}_n &= \mathbb{E}(\{T_n - \bar{T}_n\}\{T_n + \bar{T}_n\}) + \mathbb{E}(\bar{T}_n - T_n)\mathbb{E}(\bar{T}_n + T_n) \\ &\leq 2\{\mathbb{E}(\{T_n - \bar{T}_n\}^2)\mathbb{E}(\{T_n + \bar{T}_n\}^2)\}^{1/2} \leq 2\{c_3\mathbb{E}(\{T_n + \bar{T}_n\}^2)\}^{1/2} \end{aligned}$$

(c_3 is given by Lemma 4.2).

By Lemma 4.3, $\mathbb{E}(\{T_n + \bar{T}_n\}^2)$ is at most of order n^2 . Together with (4.60), this proves (1.6).

To prove (1.7), let

$$c = c(n) := c_1^2 n^{2\delta} \log n \text{ and } x_0 := n^{1+7\delta},$$

where c_1 is the constant appearing in Lemma 4.4. Then $x_0 \geq c^2$ (for large n). By Lemma 4.4 and 4.59,

$$\bar{\Delta}_k \leq c$$

and

$$\mathbb{E}(\bar{\Delta}_k^2 \mid \mathcal{F}_{k-1}) \leq \mathbb{E}(U_k \mid \mathcal{F}_{k-1}).$$

Together with Lemma 4.5, this yields (4.40), (4.41) and (4.42) (with $v = 1/2$). By applying Lemma 4.1, one obtain that for all large n and $x \leq n^{(1+7\delta)/2}$

$$\mathbb{P}(|\bar{T}_n - \mathbb{E}\bar{T}_n| > xn^{(1+7\delta)/2}) \leq c_2 e^{-c_3 x}. \quad (4.61)$$

On the other side, by Lemma 4.2, if $x > 1$ and n is large,

$$\begin{aligned} \mathbb{P}(|T_n - \mathbb{E}T_n| > xn^{(1+7\delta)/2}) &\leq \\ \mathbb{P}(|T_n - \bar{T}_n| > xn^{(1+7\delta)/2}/3) &+ \mathbb{P}(|\bar{T}_n - \mathbb{E}\bar{T}_n| > xn^{(1+7\delta)/2}/3). \end{aligned}$$

Combining this with Lemma 4.2 and (4.61) one get that for some constants $b_2, b_3 > 0$, for n large and $x \leq n^{(1+7\delta)/2}$,

$$\mathbb{P}(|T_n - \mathbb{E}T_n| > xn^{(1+7\delta)/2}) \leq b_2 e^{-b_3 x^{4\delta/(1+7\delta)}}. \quad (4.62)$$

Therefore, given $\kappa \in (1/2, 1)$ we set $\delta := (2\kappa - 1)/7$ and obtain (1.7) from (4.62).

(1.8) will follow from the same argument presented by Howard and Newman [8] to deal with the same inequality, but in the euclidean FPP context (see the proof of (4.3) there). For this reason we just sketch this proof and leave further details for the reader.

By Lemma 4.2 of Howard and Newman [8], (1.8) will follows if we prove that, for some constant c_1 ,

$$\mathbb{E}T(\mathbf{0}, 2\mathbf{n}) \geq 2\mathbb{E}T(\mathbf{0}, \mathbf{n}) - c_1 n^\kappa (\log n)^{1/\nu} \quad (4.63)$$

To prove (4.63) take $\delta < \kappa/2$ and pick $\mathbf{x}_1 := \mathbf{n}, \mathbf{x}_2, \dots, \mathbf{x}_{l(n)}$ in the boundary of D_n so that every $\mathbf{x} \in D_n$ is within distance n^δ of one of the $\mathbf{x}_j$, and $l(n) \leq cn^{1-\delta}$ for some constant $c > 0$. Let $\bar{\mathbf{x}}_j := 2\mathbf{n} - \mathbf{x}_j$ (the radial reflection of $\mathbf{x}_j$ about $\mathbf{n}$) $\bar{D}_n := 2\mathbf{n} - D_n$.

Let ρ_{2n} be a geodesic connecting $\mathbf{0}$ to $2\mathbf{n}$. Then ρ_{2n} exits D_n for the last time at some point $\mathbf{y}_n$ and after that it enters in $\bar{D}_n$ at some some point $\bar{\mathbf{y}}_n$. Let $\mathbf{x}_{j(n)} \in \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ and $\mathbf{x}_{\bar{j}(n)} \in \{\bar{\mathbf{x}}_1, \dots, \bar{\mathbf{x}}_l\}$ be the closest points to $\mathbf{y}_n$ and $\bar{\mathbf{y}}_n$ respectively.

Thus,

$$\begin{aligned} T(\mathbf{0}, 2\mathbf{n}) &\geq T(\mathbf{0}, \mathbf{y}_n) + T(\bar{\mathbf{y}}_n, 2\mathbf{n}) \\ &\geq T(\mathbf{0}, \mathbf{x}_{j(n)}) + T(\bar{\mathbf{x}}_n, 2\mathbf{n}) - [(T(\mathbf{0}, \mathbf{x}_{j(n)}) - T(\mathbf{0}, \mathbf{y}_n)) + (T(\bar{\mathbf{x}}_{j(n)}, 2\mathbf{n}) - T(\bar{\mathbf{y}}_n, 2\mathbf{n}))]. \end{aligned}$$

By sub-additivity,

$$T(\mathbf{0}, \mathbf{x}_{j(n)}) - T(\mathbf{0}, \mathbf{y}_n) \leq T(\mathbf{x}_{j(n)}, \mathbf{y}_n) \leq \max_j \sup_{\mathbf{y} - \mathbf{x}_j \leq n^\delta} T(\mathbf{x}_j, \mathbf{y})$$

and

$$T(\bar{\mathbf{x}}_{j(n)}, 2\mathbf{n}) - T(\bar{\mathbf{y}}_n, 2\mathbf{n}) \leq T(\bar{\mathbf{x}}_{j(n)}, \bar{\mathbf{y}}_n) \leq \max_j \sup_{\mathbf{y} - \mathbf{x}_j \leq n^\delta} T(\bar{\mathbf{x}}_j, \mathbf{y}).$$

Therefore,

$$\begin{aligned} T(\mathbf{0}, 2\mathbf{n}) &\geq \\ &\min_j T(\mathbf{0}, \mathbf{x}_j) + \min_j T(\mathbf{0}, \bar{\mathbf{x}}_j) - [\max_j \sup_{\mathbf{y} - \mathbf{x}_j \leq n^\delta} T(\mathbf{x}_j, \mathbf{y}) + \max_j \sup_{\mathbf{y} - \mathbf{x}_j \leq n^\delta} T(\bar{\mathbf{x}}_j, \mathbf{y})]. \end{aligned}$$

By symmetry, this implies that

$$\mathbb{E}T(\mathbf{0}, 2\mathbf{n}) \geq$$

$$2\mathbb{E}T(\mathbf{0}, \mathbf{n}) - 2[\mathbb{E}(\max_j \{\mathbb{E}T(\mathbf{0}, \mathbf{x}_j) - T(\mathbf{0}, \mathbf{x}_j)\}) - \mathbb{E}(\max_j \sup_{\mathbf{y} - \mathbf{x}_j \leq n^\delta} T(\mathbf{x}_j, \mathbf{y}))]. \quad (4.64)$$

Together with Lemma 4.3 of Howard and Newman [8], (1.7) yields

$$\mathbb{E}(\max_j \{\mathbb{E}T(\mathbf{0}, \mathbf{x}_j) - T(\mathbf{0}, \mathbf{x}_j)\}) \leq c_2 n^\kappa (\log n)^{1/\nu}. \quad (4.65)$$

By noting that the expected number of edges inside $\{\mathbf{y} : \mathbf{y} - \mathbf{x}_j \leq n^\delta\}$ is of order $n^{2\delta}$ and by using Lemma 4.3, one can prove that

$$\mathbb{E}(\max_j \sup_{\mathbf{y} - \mathbf{x}_j \leq n^\delta} T(\mathbf{x}_j, \mathbf{y})) \leq c_3 n^{2\delta} \leq c_3 n^\kappa.$$

Combining this with (4.64) and (4.65), one gets (4.63).

□

Acknowledgment. Part of this work was develop during my doctoral studies at Impa and I would like to thank my adviser, Prof. Vladas Sidoravicius, for his dedication and encouragement during this period. I also thank the whole administrative staff of IMPA for their assistance and CNPQ for financing my doctoral studies, without which this work would have not been possible. This work also amends some mistaken passages of my thesis (Pimentel [17]) and, for this reason, I should also express gratitude for the assistance given by Prof. Thomas Mountford.

REFERENCES

- [1] Alexander, K. S. (1997) Approximations of subadditive functions and convergence rates in limiting-shape results, *Ann.Probab.* **25** 30-55.
- [2] Baik, J.; Deift, P.; Johansson, K. (1999) On the distribution of the longest increasing subsequence in a random permutation, *J. Amer. Math. Soc.* **12** 1119-1178.
- [3] Cox, J. T.; Gandolfi, A.; Griffin, P. S.; Kesten, H. (1993) Greedy lattice animals I: upper bounds, *Ann. Appl. Probab.* **3**, 1151-1169.
- [4] Grimmett, G. (1999) Percolation (second edition), Springer.
- [5] Hammersley, J.M. (1961) On the rate of convergence to the connective function of the hypercubical lattice, *Quart. J. Math. Oxford, Ser. 2* **12**, 250-256.
- [6] Hammersley, J.M.; Welsh, D.J.A. (1965) First-passage percolation, sub-additive process, stochastic network and generalized renewal theory, Springer-Verlag, 61-110.
- [7] Howard, C. D.; Newman, C. M. (1997) Euclidean models of first-passage percolation, *Probab. Theory Related Fields* **108**, 153-170.
- [8] Howard, C. D.; Newman, C. M. (2001) Geodesics and spanning trees for Euclidean first-passage percolation, *Ann. Probab.* **29**, 577-623.
- [9] Johansson, K. (2000) Shape fluctuations and random matrices, *Comm. Math. Phys.* **209**, 437-476.
- [10] Kardar, K.; Parisi, G.; Zhang Y. Z. (1986) Dynamic scaling of growing interfaces *Phys. Rev. Lett.* **56**, 889-892.
- [11] Kesten, H. (1963) On the number of self-avoiding walks *J. Math. Phys.* **4**, 960-969.
- [12] Kesten, H. (1986) Aspects of first-passage percolation, *Lectures Notes in Math.* **1180**, Springer-Verlag, 125-264
- [13] Kesten, H. (1993) On the speed of convergence in first-passage percolation *Ann. Appl. Probab.* **3**, 296-338.
- [14] Krug, J.; Spohn, H. (1991) *Solids far from Equilibrium*, (C. Godreche, ed.) Cambridge Univ. Press, Cambridge.
- [15] Ligget, T.M.; Schonmann, R.H.; Stacey, A.M. (1997) Domination by product measures, *Ann. Probab.* **25**, 71-95.
- [16] Moller, J. (1991) Lectures on random Voronoi tessellations, *Lectures Notes in Stat.* **87**, Springer-Verlag.

- [17] Pimentel, L. P. R. (2004) Competing growth, interfaces and geodesics in first-passage percolation on Voronoi tilings. Phd Thesis, IMPA, Rio de Janeiro.
- [18] Talagrand, M. (1995), Concentration of measure and isoperimetric inequalities in product spaces *Inst. Hautes Etudes Sci. Publ. Math.* **81**, 73-205.
- [19] Vahidi-Asl, M.Q.; Wierman, J.C. (1990) First-passage percolation on the Voronoi tessellation and Delaunay triangulation, Random Graphs 87 (M. Karonske, J. Jaworski and A. Rucinski, eds.), 1990, 341-359.
- [20] Vahidi-Asl, M.C.; Wierman, J.C. (1992) A shape result for first-passage percolation on the Voronoi tessellation and Delaunay triangulation, Random Graphs 89 (A. Frieze and T. Luczak, eds.), Wiley, 247-262.

INSTITUT DE MATHÉMATIQUES, ÉCOLE POLYTECHNIQUE FÉDÉRALE DE LAUSANNE, CH-1015 LAUSANNE, SWITZERLAND.

E-mail address: leandro.pimentel@epfl.ch

URL: <http://ima.epfl.ch/~lpimente/>